

Slip acceleration generates seismic tremor like signals in friction experiments

Dimitri Zigone,¹ Christophe Voisin,^{2,3} Eric Larose,¹ François Renard,^{4,5} and Michel Campillo¹

Received 24 September 2010; revised 4 November 2010; accepted 15 November 2010; published 15 January 2011.

[1] Since their discovery nearly a decade ago, the origin of seismic tremor remains unclear. Recent studies indicate that various driving phenomena such as Earth and ocean tides, regional and teleseismic earthquakes enhance tremor activity. Observations of the coincidence with slow-slip events and of fast migrations of tremors have led frictional slip to be considered as the possible source of tremors. Indeed, laboratory friction experiments succeeded in generating and recording tremor like signals (TLS). Here we show a systematic correlation between the onset of slip acceleration and the emission of TLS in a laboratory friction experiment. TLS are generated when the shear stress reaches the peak static resistance and the dilatancy meets its maximum that is when the mature interface is close to failure. This robust result provides a comprehensive image of how natural seismic tremors might be generated and/or triggered by passing seismic waves, tides or even slow slip events. **Citation:** Zigone, D., C. Voisin, E. Larose, F. Renard, and M. Campillo (2011), Slip acceleration generates seismic tremor like signals in friction experiments, *Geophys. Res. Lett.*, 38, L01315, doi:10.1029/2010GL045603.

1. Introduction

[2] Seismic tremors, named also non-volcanic tremors, have now been well documented and studied in many subduction zones [Dragert *et al.*, 2004; Hirose and Obara, 2006; Kao *et al.*, 2005; Obara, 2002; Obara and Hirose, 2006; Shelly *et al.*, 2007b, 2006] and along some continental fault segments [Ghosh *et al.*, 2009b; Nadeau and Dolenc, 2005; Peng *et al.*, 2009, 2008]. Recent studies indicate that various driving phenomena such as Earth and ocean tides [Thomas *et al.*, 2009], regional and teleseismic earthquakes [Ghosh *et al.*, 2009b; Peng *et al.*, 2009, 2008; Rubinstein *et al.*, 2009] enhance tremor activity. Frictional slip was proposed as the possible source of tremors [Brown *et al.*, 2009; Ghosh *et al.*, 2009a; Ide *et al.*, 2007; Kao *et al.*, 2007; La Rocca *et al.*, 2009; Larmat *et al.*, 2009].

[3] Laboratory experiments specifically designed to study seismic tremors are a powerful tool to explore the physical processes at the origin of these signals. Few experiments

have been successful in producing tremor like signals (TLS) so far. They involve fluid-flow and fluid processes [Burlini *et al.*, 2009], or frictional processes associated with shear of a deformable sample [Voisin *et al.*, 2007, 2008]. Our experimental setup (Figure 1a) is designed to reproduce different frictional behaviors, from stick-slip to stable sliding. We make use of a deformable slider of salt (NaCl) pushed at constant load-point velocity, under constant conditions of normal pressure (0.26 MPa), temperature (22°C) and ambient humidity. Using a salt slider allows for the brittle and ductile deformation to be effective on the time scale of our experiments, aimed to serve as an analogue for natural faults deforming in the brittle and ductile regimes at 20–40 km depth. We continuously record the frictional force and the acoustic emissions generated during the shear of the sample. These signals are carefully scrutinized in order to investigate the temporal timing of the sliding characteristics and of the TLS emission.

2. Results

[4] Figures 1b–1d present three different stages of a friction experiment representative of the stable, intermediate and unstable frictional regimes. The unstable regime is characterized by cycles of long stress increase followed by sudden drops of the friction force, associated with the stick-slip behavior of the slider (stage 1, Figure 1b). The jumps of the slider produce short duration and high amplitude acoustic signals. In this stage of stick-slip, no TLS is recorded. The salt slider follows an evolution from an unstable stick-slip behavior to more and more stable behavior with accumulated displacement [Voisin *et al.*, 2007]. Figure 1c presents a second experiment (under the same conditions) where the sample has accumulated 3.3 mm of displacement and for which the slider interface has become more mature (stage 2). In this condition, the stick-slip behavior changes and smoothens. The slider still obeys a stick-slip behavior, but is creeping before and after the jump. Interestingly, the associated acoustic emission presents two types of signals: i) a tremor like signal with long duration and low amplitude followed by ii) a strong impulsive and short duration event, signature of the jump. Finally, Figure 1d presents an experiment in the stable regime, characterized by smooth oscillations of the frictional force around a mean value, which corresponds to smooth variations of the sliding velocity of the slider [Voisin *et al.*, 2007]. During this stable regime, the acoustic emission is composed exclusively of noise and TLS. Those TLS are identical to those presented in Figure 1c in terms of frequency content, duration and amplitude, suggesting their common physical origin. These results show that TLS in

¹Laboratoire de Géophysique Interne et Tectonophysique, CNRS, Université Joseph Fourier, Grenoble, France.

²Laboratoire de Géophysique Interne et Tectonophysique, IRD, CNRS, Université Joseph Fourier, Grenoble, France.

³Centre de Recherches Géophysiques, CNRSL, Bhanès, Liban.

⁴Laboratoire de Géodynamique des Chaînes Alpines, CNRS, Université Joseph Fourier, Grenoble, France.

⁵Physics of Geological Processes, University of Oslo, Oslo, Norway.




Figure 1. (a) Sketch of the experimental block-slider system (details in the auxiliary materials).¹ Frictional behavior of a salt/glass interface (top plots: recorded shear force divided by normal load, noted F/W) and associated acoustic emission records (bottom plots). (b) Typical stick-slip regime characterized by sudden shear stress drops separated by periods of stress accumulation. The acoustic emission is formed of large impulsive events, sometimes preceded and followed by smaller impulsive events. (c) The frictional behavior is more complex than previously. If the sudden jumps still occur, they are preceded by smooth oscillations of growing amplitude. This behavior is at the limit between stable and unstable behaviors. It arises with the cumulative displacement of the slider (here 3.3 mm; about 120 cycles) that modifies the properties of the salt interface. The associated acoustic emission presents a complex signal formed by a TLS with long duration (10 to 20 ms) and low amplitude followed by a strong impulsive and short duration event that represents the signature of the jump. (d) The frictional force remains more or less constant with small variations around a mean value (see insert): this is the stable regime obtained after 8mm (about 150 cycles) of cumulative slip. The associated acoustic emission is formed of a TLS with low amplitude and long duration, emitted at each slip acceleration, when the shear stress and the dilatancy are at maximum.

friction experiments are related to regimes of stable slip (Figure 1d) or simultaneous creep and stick-slip (Figure 1c). Note however, that the high level of experimental noise impairs to exclude definitively the existence of TLS at other time than the slip events.

[5] Figure 2 presents 4 typical examples out of the 46 TLS extracted from a friction experiment (see Figure S1 of the auxiliary materials for an exhaustive presentation of the TLS).¹ For each case, the TLS slowly emerges from the background noise, keeping a low waxing and waning amplitude. Some

bursts or peaks in the signal occur randomly, creating some variability in the TLS. The maximum amplitude of TLS is always relatively small, in the range $\pm 5 \cdot 10^{-7} \text{ m} \cdot \text{s}^{-1}$. The apparent duration of a TLS is variable, with 90% of the records in the range 8–20 milliseconds. We compared the Fourier power spectrum of the acoustic events recorded during the experiments against the Fourier spectrum of background noise (Figure 3). The acoustic event associated with a stick-slip event (see Figure 1b for the time plot) presents a highly energetic spectrum with a few peaks between 10^4 and 10^5 Hertz. Conversely, the TLS spectrum is hardly above the noise spectrum in this same frequency range, confirming the low S/N ratio of these signals. Any-

¹Auxiliary materials are available in the HTML. doi:10.1029/2010GL045603.




Figure 2. Examples of TLS and the associated slip acceleration evolution. We present here 4 examples out of the 46 windows containing TLS (see Figure S1 in auxiliary materials). All TLS share some common features: progressive emergence from the background noise, very low amplitude, and rather long duration of 10–20 ms. Here and there, some bursts of energy create a variability of the TLS. Associated with the time traces of the TLS, we represent the shear stress evolution in a manner that mimics the slip acceleration. The high sampling rate of both force and acoustic emission and the short instrumental response of a millisecond allow for a precise comparison between acceleration of slip (top plot) and acoustic record (bottom plot). For each event A, B, C, and D, an arrow points to the beginning of slip acceleration determined by visual inspection. We note the strong correlation between the onset of slip acceleration and the emission of the TLS. The maximum of the TLS is reached after a few milliseconds. Note all TLS do occur at the beginning of the acceleration, and are thus associated with change of slip properties of the frictional interface.

way, we can observe two peaks of energy at 48 and 57 kHz. The strong resonance inherent to the experimental set-up and the limited bandwidth of the sensor whose response is flat in the 10–60 kHz range probably alter the shape of the TLS spectrum. Nonetheless, most of the energy of the TLS apparently lies in the range 40–60 kHz. Such a frequency range corresponds to a wavelength of a few centimeters and indicates vibrations of the whole NaCl sample, and impairs the description of the microstructural process(es) at work during the emission of the TLS.

[6] A continuous record of the acoustic emission together with the frictional force at a 500 kHz sampling frequency allows for a precise timing of the TLS with respect to the onset of slip acceleration. Doing such a careful analysis for all the recorded TLS we observe a systematic correlation between the onset of slip acceleration of the interface and the generation of TLS. Figure 2 exemplifies this behavior where the acceleration of slip (top plot) is associated with a burst of acoustic emissions (bottom plot) for the 4 selected slip events. There is a clear temporal correlation between the

occurrence of TLS and the beginning of slip acceleration that starts when the shear stress equals the static resistance and the maximum level of dilatancy sustained by the slider is reached. The relation between shear stress level, dilatancy and tremor like signals might explain why the TLS progressively vanish while the slider is still accelerating. We might infer from the experiments that the sliding of the interface is associated with a change in the contacts population resulting from the mass redistribution at the interface and leading to the generation of a TLS right before the drop in friction.

[7] However, if slip acceleration is a necessary condition, it is not the only one to be met to emit a TLS. It is interesting to remark that TLS are not recorded at the beginning of the experiment, during the stick-slip stage (Figure 1b). The interface has to accumulate some amount of slip before to emit TLS. This might be related to the restructuring of the sliding interface and the development of the striations parallel to the sliding direction [Voisin *et al.*, 2007, 2008]. A third condition has to be met, related to the state of stress




Figure 3. Fourier spectrums of the acoustic signature of a jump (blue curve), TLS (red curve), and noise (black curve) computed from 20 ms time windows. The low frequency range of the signals recorded below 10 kHz is not presented on the figure because it is dominated by experimental noise (motor) and associated with the movement of the slider as a block. The noise spectrum is computed from the mean of different time windows. The spectrum of a jump is highly energetic between 10 kHz and 100 kHz. By contrast the spectrum of the TLS is close to the noise level and presents only two clear peaks of energy at 48 and 57 kHz, not present in the power spectrum of the jump signal.

and/or the peak dilatancy of the slider. TLS are not recorded randomly. During stage 2 (Figure 1c), TLS are recorded only during the last smooth oscillation before the stick-slip, that is when the interface is close to failure. During stage 3 (Figure 1d), TLS are recorded at each slip acceleration that occurs when the shear stress is at a maximum and starts to decrease gently. In both cases, the TLS occur at the maximum of the shear stress and also at the maximum of dilatancy bore by the slider.

3. Discussion and Conclusions

[8] TLS recorded in these experiments are related solely to the frictional process occurring along the contact interface. Because of the limited wavelength of the sensors used in this study, we are not able to characterize the processes at the origin of the TLS. Consequently, there is no simple possibility to derive a scaling between TLS and natural seismic tremors. Nonetheless, if we assume that the same process could produce seismic tremors in the nature as suggested by the coincidence of tremors with slow-slip events or by the fast migrations of tremors [Brown *et al.*, 2009; Ghosh *et al.*, 2009a; Ide *et al.*, 2007; Kao *et al.*, 2007; La Rocca *et al.*, 2009; Larmat *et al.*, 2009], we can use the laboratory results to shed some light on the natural system of slow slip, seismic tremors and triggering.

[9] Result 1: This friction experiment conducted with a deformable interface exhibits a large variety of frictional behavior associated with different amounts of cumulative slip. These behaviors are themselves accompanied by different acoustic signals, impulsive events and TLS. A comparable variety of behaviors is observed on natural fault segments exhibiting in time and space different frictional

behaviors: seismic, aseismic, creeping, associated with seismic events or seismic tremors.

[10] Result 2: the TLS are emitted when the shear stress and/or the dilatancy are at maximum. A large and growing number of observations emphasize the link between stress changes and seismic tremors. Maybe the clearest evidence is the triggering of seismic tremor by large transient shear stresses [Rubinstein *et al.*, 2007], dilatational stresses [Miyazawa and Mori, 2006], or the tidal modulation of tremor rate [Rubinstein *et al.*, 2008]. Such conveniently orientated stress waves temporarily increase the stress level on a given sliding interface or increase the dilatancy, inducing slip and tremor triggering like in the laboratory experiments. Another evidence might be found in silent or slow slip events that are often associated with seismic tremors in episodic tremor and slip events, described for the first time in Cascadia [Dragert *et al.*, 2001]. Tremors are sometimes found to migrate along strike, in agreement with the slow slip propagation [Obara and Sekine, 2009; Shelly *et al.*, 2007a]. Lead by the laboratory results, we propose that the temporary increase of stress induced by the rupture front propagation itself can trigger seismic tremors in zones where the stress state is close to its maximum. The present understanding of rupture propagation relies on a friction law, either slip-dependent or rate-and-state dependent, that defines a breaking-down zone governing the stress drop preceded by a short stress increase [Ida, 1975; Rubin, 2008; Voisin *et al.*, 2002]. Assuming that a similar breaking down process occurs for slow slip event, the small stress increase associated with the rupture tip would be able to trigger seismic tremors, according to our experimental results.

[11] Result 3: TLS occur when the slider accelerates. Seismic tremors appear as a local recollection of the unstable frictional behavior occurring during the slippage of aseismic slow events down dip subduction zones [Shelly *et al.*, 2007a]. At a large scale, a slow slip event can be considered as an acceleration of slip on the interface during a few months, thus a possible source for seismic tremors. The slip complexity derived from inversion imposes slip rate variations and thus local accelerations that would be able to trigger seismic tremors.

[12] Result 4: TLS are emitted only when enough slip is accumulated and the interface has developed a striation [Voisin *et al.*, 2007]. It is the case down-dip the subduction zones, where seismic tremors were first recorded. The recent discovery of the control of tremor migration by preferred directions linked to striation of the subduction plane [Ide, 2010] together with our experimental observations suggest that the striation is a necessary condition to emit seismic tremors and to control the tremor migration [Ide, 2010]. Large cumulative slips are also observed along some large continental strike slip faults. The relation between seismic tremors and seismic events in continental context remains open. If most of the seismic tremors and seismic events occur in different areas, the brittle crust for the latter, and the deeper ductile crust for the former, in very rare cases they do occur beneath the same location. It is the case with the Cholame area, a segment of the San Andreas Fault at the north tip of the great 1857 Fort Tejon earthquake rupture. Tremor activity was detected just before and after the 2004 M6 Parkfield earthquake, at the depth where the earthquake rupture nucleated, at the transition between the stable sliding layer and the seismogenic layer [Nadeau and Guilhem,

2009]. Our experiment reported in Figure 1c shows that the same interface can generate tremors and seismic events. We observe that the TLS emission occurs systematically before the slip event. Because of the complete stress release in our experiment, the TLS rapidly stops after a few milliseconds. This is to be compared to the Cholame segment, where the “fore-tremor” activity was proposed before to the Parkfield earthquake of 2004. A recent study (M. Bouchon et al., Observation of tremors before a large earthquake, submitted to *Science*, 2010) also reports LFEs and tremors located in the hypocentral area before the large 1999 Izmit earthquake. We thus can imagine that the tremors would be generated by some creep acceleration eventually leading to the seismic event [Shelly, 2009]. If so, seismic tremors occurring at the base of the brittle crust might be seen as potential signature of the nucleation of seismic events.

[13] **Acknowledgments.** This study was supported by ANR-09-JCJC-0011-01 grant. We acknowledge the technical support of Sophie Beauprêtre, Raphaël Jacquot, Adeline Richard, Sandrine Roussel, Benjamin Vial, and Liliane Jenaton. The authors thank Ruth Harris, André Niemeijer and an anonymous reviewer for their constructive reviews.

References

- Brown, J. R., G. C. Beroza, S. Ide, K. Ohta, D. R. Shelly, S. Y. Schwartz, W. Rabbel, M. Thorwart, and H. Kao (2009), Deep low-frequency earthquakes in tremor localize to the plate interface in multiple subduction zones, *Geophys. Res. Lett.*, *36*, L19306, doi:10.1029/2009GL040027.
- Burlini, L., G. Di Toro, and P. Meredith (2009), Seismic tremor in subduction zones: Rock physics evidence, *Geophys. Res. Lett.*, *36*, L08305, doi:10.1029/2009GL037735.
- Dragert, H., K. L. Wang, and T. S. James (2001), A silent slip event on the deeper Cascadia subduction interface, *Science*, *292*(5521), 1525–1528, doi:10.1126/science.1060152.
- Dragert, H., K. Wang, and G. Rogers (2004), Geodetic and seismic signatures of episodic tremor and slip in the northern Cascadia subduction zone, *Earth Planets Space*, *56*(12), 1143–1150.
- Ghosh, A., J. E. Vidale, J. R. Sweet, K. C. Creager, and A. G. Wech (2009a), Tremor patches in Cascadia revealed by seismic array analysis, *Geophys. Res. Lett.*, *36*, L17316, doi:10.1029/2009GL039080.
- Ghosh, A., J. E. Vidale, Z. Peng, K. C. Creager, and H. Houston (2009b), Complex nonvolcanic tremor near Parkfield, California, triggered by the great 2004 Sumatra earthquake, *J. Geophys. Res.*, *114*, B00A15, doi:10.1029/2008JB006062.
- Hirose, H., and K. Obara (2006), Short-term slow slip and correlated tremor episodes in the Tokai region, central Japan, *Geophys. Res. Lett.*, *33*, L17311, doi:10.1029/2006GL026579.
- Ida, Y. (1975), Analysis of stick-slip and earthquake mechanism, *Phys. Earth Planet. Inter.*, *11*(2), 147–156, doi:10.1016/0031-9201(75)90008-4.
- Ide, S. (2010), Striations, duration, migration and tidal response in deep tremor, *Nature*, *466*(7304), 356–359, doi:10.1038/nature09251.
- Ide, S., D. R. Shelly, and G. C. Beroza (2007), Mechanism of deep low frequency earthquakes: Further evidence that deep non-volcanic tremor is generated by shear slip on the plate interface, *Geophys. Res. Lett.*, *34*, L03308, doi:10.1029/2006GL028890.
- Kao, H., S. J. Shan, H. Dragert, G. Rogers, J. F. Cassidy, and K. Ramachandran (2005), A wide depth distribution of seismic tremors along the northern Cascadia margin, *Nature*, *436*(7052), 841–844, doi:10.1038/nature03903.
- Kao, H., S.-J. Shan, G. Rogers, and H. Dragert (2007), Migration characteristics of seismic tremors in the northern Cascadia margin, *Geophys. Res. Lett.*, *34*, L03304, doi:10.1029/2006GL028430.
- Larmat, C. S., R. A. Guyer, and P. A. Johnson (2009), Tremor source location using time reversal: Selecting the appropriate imaging field, *Geophys. Res. Lett.*, *36*, L22304, doi:10.1029/2009GL040099.
- La Rocca, M., K. C. Creager, D. Galluzzo, S. Malone, J. E. Vidale, J. R. Sweet, and A. G. Wech (2009), Cascadia tremor located near plate interface constrained by S minus P wave times, *Science*, *323*(5914), 620–623, doi:10.1126/science.1167112.
- Miyazawa, M., and J. Mori (2006), Evidence suggesting fluid flow beneath Japan due to periodic seismic triggering from the 2004 Sumatra-Andaman earthquake, *Geophys. Res. Lett.*, *33*, L05303, doi:10.1029/2005GL025087.
- Nadeau, R. M., and D. Dolenc (2005), Nonvolcanic tremors deep beneath the San Andreas Fault, *Science*, *307*(5708), 389, doi:10.1126/science.1107142.
- Nadeau, R. M., and A. Guilhem (2009), Nonvolcanic tremor evolution and the San Simeon and Parkfield, California, earthquakes, *Science*, *325*(5937), 191–193, doi:10.1126/science.1174155.
- Obara, K. (2002), Nonvolcanic deep tremor associated with subduction in southwest Japan, *Science*, *296*(5573), 1679–1681, doi:10.1126/science.1070378.
- Obara, K., and H. Hirose (2006), Non-volcanic deep low-frequency tremors accompanying slow slips in the southwest Japan subduction zone, *Tectonophysics*, *417*(1–2), 33–51, doi:10.1016/j.tecto.2005.04.013.
- Obara, K., and S. Sekine (2009), Characteristic activity and migration of episodic tremor and slow-slip events in central Japan, *Earth Planets Space*, *61*(7), 853–862.
- Peng, Z., J. E. Vidale, K. C. Creager, J. L. Rubinstein, J. Gomberg, and P. Bodin (2008), Strong tremor near Parkfield, CA, excited by the 2002 Denali Fault earthquake, *Geophys. Res. Lett.*, *35*, L23305, doi:10.1029/2008GL036080.
- Peng, Z., J. E. Vidale, A. G. Wech, R. M. Nadeau, and K. C. Creager (2009), Remote triggering of tremor along the San Andreas Fault in central California, *J. Geophys. Res.*, *114*, B00A06, doi:10.1029/2008JB006049.
- Rubin, A. M. (2008), Episodic slow slip events and rate-and-state friction, *J. Geophys. Res.*, *113*, B11414, doi:10.1029/2008JB005642.
- Rubinstein, J. L., J. E. Vidale, J. Gomberg, P. Bodin, K. C. Creager, and S. D. Malone (2007), Non-volcanic tremor driven by large transient shear stresses, *Nature*, *448*(7153), 579–582, doi:10.1038/nature06017.
- Rubinstein, J. L., M. La Rocca, J. E. Vidale, K. C. Creager, and A. G. Wech (2008), Tidal modulation of nonvolcanic tremor, *Science*, *319*(5860), 186–189, doi:10.1126/science.1150558.
- Rubinstein, J. L., J. Gomberg, J. E. Vidale, A. G. Wech, H. Kao, K. C. Creager, and G. Rogers (2009), Seismic wave triggering of nonvolcanic tremor, episodic tremor and slip, and earthquakes on Vancouver Island, *J. Geophys. Res.*, *114*, B00A01, doi:10.1029/2008JB005875.
- Shelly, D. R. (2009), Possible deep fault slip preceding the 2004 Parkfield earthquake, inferred from detailed observations of tectonic tremor, *Geophys. Res. Lett.*, *36*, L17318, doi:10.1029/2009GL039589.
- Shelly, D. R., G. C. Beroza, S. Ide, and S. Nakamura (2006), Low-frequency earthquakes in Shikoku, Japan, and their relationship to episodic tremor and slip, *Nature*, *442*(7099), 188–191, doi:10.1038/nature04931.
- Shelly, D. R., G. C. Beroza, and S. Ide (2007a), Complex evolution of transient slip derived from precise tremor locations in western Shikoku, Japan, *Geochem. Geophys. Geosyst.*, *8*, Q10014, doi:10.1029/2007GC001640.
- Shelly, D. R., G. C. Beroza, and S. Ide (2007b), Non-volcanic tremor and low-frequency earthquake swarms, *Nature*, *446*(7133), 305–307, doi:10.1038/nature05666.
- Thomas, A. M., R. M. Nadeau, and R. Burgmann (2009), Tremor-tide correlations and near-lithostatic pore pressure on the deep San Andreas fault, *Nature*, *462*(7276), 1048–1051, doi:10.1038/nature08654.
- Voisin, C., I. Ionescu, and M. Campillo (2002), Crack growth resistance and dynamic rupture arrest under slip dependent friction, *Phys. Earth Planet. Inter.*, *131*(3–4), 279–294, doi:10.1016/S0031-9201(02)00054-7.
- Voisin, C., F. Renard, and J.-R. Grasso (2007), Long term friction: From stick-slip to stable sliding, *Geophys. Res. Lett.*, *34*, L13301, doi:10.1029/2007GL029715.
- Voisin, C., J.-R. Grasso, E. Larose, and F. Renard (2008), Evolution of seismic signals and slip patterns along subduction zones: Insights from a friction lab scale experiment, *Geophys. Res. Lett.*, *35*, L08302, doi:10.1029/2008GL033356.

M. Campillo, E. Larose, and D. Zigone, Laboratoire de Géophysique Interne et Tectonophysique, CNRS, Université Joseph Fourier, Maison des Géosciences, BP 53, F-38041 Grenoble CEDEX, France. (dimitri.zigone@ujf-grenoble.fr)

F. Renard, Laboratoire de Géodynamique des Chaînes Alpines, CNRS, Université Joseph Fourier, Maison des Géosciences, BP 53, F-38041 Grenoble CEDEX, France.

C. Voisin, Laboratoire de Géophysique Interne et Tectonophysique, IRD, CNRS, Université Joseph Fourier, Maison des Géosciences, BP 53, F-38041 Grenoble CEDEX, France.